MAKING A DIFFERENCE IN THE LIVES OF CHILDREN, YOUTH AND FAMILIES



## Illness Management- Stress dosing for Adrenal Insufficiency

Division of Endocrinology and Metabolism



## ILLNESS MANAGEMENT: STRESS DOSING FOR ADRENAL INSUFFICIENCY

## **Medication Dose**

For fever between 37.5 - 38.5 C orally or underarm (38-39 rectal) and /or sore throat and /or cough and or minor injury <a href="Mountainser">DOUBLE or increase</a> the doses to

AM\_\_\_\_\_\_

\_\_\_\_\_

PM\_\_\_\_\_

Bedtime\_

Affix Label

## **Emergency Care**

You may have been taught to give the medication by injection (Solu-Cortef). This is required if your child:

- is unable to keep down the medication by mouth
- is unable to respond normally, is unconscious or if they have had a serious injury.

\*\*You may also need to call 911\*\*

Call the <u>Endocrinologist-on-call</u> for directions and prepare to go to the nearest Emergency department for assessment, possible IV medication and hydration.

- Present your wallet card to the triage nurse and inform them that your child has
- Adrenal insufficiency /Addison's disease /CAH

The Emergency Department will contact the **Endocrinologist-on-call**.

All children with permanent forms of adrenal insufficiency require a <u>medic alert bracelet</u>. Parents of children with Adrenal Suppression may discuss this need with their endocrinologist.

For fever greater than 38.5 C orally or underarm (39 rectal), vomiting and/or diarrhea or a severe injury TRIPLE or increase the doses to

AM\_\_\_\_\_

PM\_\_\_\_\_

Bedtime\_\_\_\_\_

You can call the Endocrinologist-on-call during illness/injury with any urgent concerns about the above instructions.

CHEO 613-737-7600 x 0 and ask for the Endocrinologist–on-call

You may also call your Endocrine Nurse during business hours for non-urgent concerns 613-737-7600 ext

RN:\_\_\_\_